arXiv:nucl-th/9710014v2 11 May 1998

Thermodynamics of Δ resonances

W. Weinhold[1], B. Friman, W. Nörenberg

Gesellschaft für Schwerionenforschung (GSI),
Planckstr. 1, D-64291 Darmstadt, Germany

and

Institut für Kernphysik, Technische Universität Darmstadt,
Schloßgartenstr. 9, D-64289 Darmstadt, Germany

Abstract

The thermodynamic potential of a system of pions and nucleons is computed including the πN interactions in the P_{33} channel. A consistent treatment of the width of the resonance in this channel, the $\Delta(1232)$ resonance, is explored in detail. In the low-density limit we recover the leading term of the virial expansion for the thermodynamic potential. An instructive diagrammatic interpretation of the contributions to the total baryon number is presented. Furthermore, we examine within a fireball model the consequences for the pion spectra in heavy-ion collisions at intermediate energies, including the effect of collective flow. A consistent treatment of the Δ width leads to a substantial enhancement of the pion yield at low momenta.

---

1 Introduction

In heavy-ion collisions a zone of hot and dense hadronic matter is created. At beam energies on the order of 1 GeV per nucleon or higher, hadron resonances play an important role in the dynamics of such collisions. Some of these resonances have large widths, comparable to or larger than the typical temperature of the interaction zone. Also stable hadrons generally acquire a width in dense matter, due to the interactions with the medium. Consequently, a consistent theory of the thermodynamics and kinetics of many-body systems with short-lived particles is necessarily a basic element in the description of such collisions. At present a satisfactory treatment of this problem is still lacking.

In this letter we explore the consequences of the width of the $\Delta(1232)$ resonance on the thermodynamics of hot and dense hadronic matter. In heavy-ion

---

[1] e-mail: w.weinhold@gsi.de

Preprint submitted to Physics Letters B *6 February 2008*

collisions at beam energies around 1 GeV per nucleon, the nucleon and the pion are together with the $\Delta(1232)$ the most abundant hadrons. By treating the $\Delta(1232)$ as an independent field, one effectively accounts for the pion-nucleon interactions in the resonant P_{33} channel. As a first approximation we assume that the interaction zone consists only of these particles. The extension of our approach to other channels is straightforward.

The thermodynamics of such systems has been studied by many authors. In calculations based on the Green-function or related techniques the width of the Δ was, so far, either neglected or dealt with in an *ad hoc* manner [1–4]. On the other hand it is well known how to take interactions in general and resonances in particular into account within the S-matrix formulation of statistical mechanics [5,6]. In section 2 we illustrate the relation between the two techniques within a model.

In order to assess the consequences of the Δ width for the pion spectrum in heavy-ion collisions we consider a simple fireball model (section 3). In this model it is assumed that the interaction region, created in such collisions, is in local thermal and chemical equilibrium at freeze-out, i.e., when the particles decouple. There are two sources for the observed pions: the free pions with a thermal distribution present at freeze-out and the pions emanating from the decay of thermally excited Δ-resonances. We show that popular prescriptions for evaluating the resonance contribution are incorrect.

2 Thermodynamic potential and baryon density

Our starting point for computing the thermodynamic potential Ω of a πNΔ-gas as a function of temperature T, baryon chemical potential μ_B and volume V is the Green-function approach of Baym and Kadanoff [7]. The temperature Green functions are denoted by $\mathcal{G}_N$, $\mathcal{G}_\Delta$ for the nucleon and Δ resonance respectively, and by $\mathcal{D}_\pi$ for the pion. The free Green functions are defined by

$$\mathcal{G}^0_B(\vec{p}, E_m) = \left[(iE_m - \mu_B) - \epsilon_B(\vec{p})\right]^{-1}, \qquad \epsilon_B(\vec{p}) = m^0_B + \vec{p}^{\,2}/2m^0_B \tag{1}$$

$$\mathcal{D}^0_\pi(\vec{q}, \omega_m) = \left[(i\omega_m)^2 - (\omega_\pi(\vec{q}))^2\right]^{-1}, \qquad \omega_\pi(\vec{q}) = \sqrt{\vec{q}^{\,2} + m^{0\,2}_\pi}\,, \tag{2}$$

where $B = N, \Delta$ while E_m and ω_m denote the Matsubara frequencies for fermions and bosons, respectively. We consider an interaction vertex of the $\pi N\Delta$ type, which connects a pion and a nucleon with a Δ resonance. The following discussion depends only on the topology of the diagrams considered and not on a particular choice of the coupling. Hence, we specify the form of the interaction only at the end of this section, where numerical results are presented.



Fig. 1. Diagrammatic representation of (a) the functional Φ and (b) contributions to the baryon density. A double line denotes a Δ, a solid line a nucleon and a dashed line a pion Green function. Thick lines correspond to dressed, thin lines to free Green functions. The gray circle indicates the baryon number operator. The first diagram in (b) yields the nucleon density n_N, the second one the density of Δ resonances in the system n_Δ, while the third one, δn_N, corresponds to the πN component of a physical Δ (see eqs. (8) and (9)).

The exact thermodynamic potential Ω of an interacting system of pions, nucleons and Δ resonances can be expressed in terms of the full Green functions [7–9]

$$\begin{aligned}\Omega = &\frac{1}{2}\,\mathrm{Tr}\,\ln\left[-\mathcal{D}_\pi^{-1}\right] + \frac{1}{2}\,\mathrm{Tr}\,\Pi_\pi\mathcal{D}_\pi - \mathrm{Tr}\,\ln\left[-\mathcal{G}_N^{-1}\right] - \mathrm{Tr}\,\Sigma_N\mathcal{G}_N \\ &-\mathrm{Tr}\,\ln\left[-\mathcal{G}_\Delta^{-1}\right] - \mathrm{Tr}\,\Sigma_\Delta\mathcal{G}_\Delta + \Phi\,, \end{aligned} \tag{3}$$

where Σ_N, Σ_Δ and Π_π are the self energies of the nucleon, the Δ resonance and the pion. The traces correspond to sums over spin, isospin, and Matsubara frequencies as well as integrations over phase space. The functional Φ is the sum of all two-line irreducible skeleton diagrams where the lines represent full Green functions. The self energy of a particle is obtained by a functional variation of Φ with respect to the Green function: $\Sigma_B = \delta\Phi/\delta\mathcal{G}_B$ and $\Pi_\pi = -2\ \delta\Phi/\delta\mathcal{D}_\pi$. Diagrammatically, the functional variation is accomplished by opening the corresponding line [8].

As shown by Baym and Kadanoff [7], the choice of a subset of diagrams for Φ defines a self-consistent approximation, where every Green function is dressed by its self energy obtained by the variation of Φ. The advantage of such so-called conserving or Φ-derivable approximations is that they satisfy conservation laws and guarantee thermodynamic consistency. This approach has in the past been applied to a wide variety of many-body problems [9]. For the πNΔ system the simplest approximation of this type is obtained by choosing Φ equal to the two-loop diagram shown in fig. 1(a). The corresponding self energy of the Δ resonance is obtained by opening the Δ line.

A fully self-consistent treatment of this problem would be quite involved, and outside the scope of this exploratory calculation. We therefore relax the self-consistency and consider a low-density expansion[2] of eq. (3). In the pion and nucleon Green functions we retain only the pole terms, i.e., we use eqs. (1-2), albeit with physical masses. On the other hand, for the Δ, which has a large width, the vacuum self energy Σ_Δ is treated explicitly. One then finds, to leading order in the baryon density

$$\Omega = \frac{1}{2}\,\mathrm{Tr}\,\ln\Big[-\left(\mathcal{D}_\pi^V\right)^{-1}\Big] - \mathrm{Tr}\,\ln\Big[-\left(\mathcal{G}_N^V\right)^{-1}\Big] - \mathrm{Tr}\,\ln\Big[-\left(\mathcal{G}_\Delta^V\right)^{-1}\Big]\,, \qquad (4)$$

where $\mathcal{D}_\pi^V = \mathcal{D}_\pi^0$ and $\mathcal{G}_N^V = \mathcal{G}_N^0$ with $m_\pi^0 = m_\pi$ and $m_N^0 = m_N$, respectively and $\left(\mathcal{G}_\Delta^V\right)^{-1} = \left(\mathcal{G}_\Delta^0\right)^{-1} - \Sigma_\Delta$. Since the thermodynamic potential in eq. (4) is a sum of independent terms involving the vacuum Green functions, it may superficially appear that the problem has now been reduced to a trivial one with non-interacting pions, nucleons and Δ resonances. However, as we show below, this approximation includes non-trivial interaction effects. In fact, the Δ-resonance term accounts for the πN interactions in the P_{33} channel.

The traces in eq. (4) can be evaluated using standard techniques (cf. [12]). For a spin and isospin symmetric system one finds

$$\begin{aligned}\Omega\left(T,\mu_B,V\right) = {} & 3\,TV\int\frac{\mathrm{d}^3q}{\left(2\pi\right)^3}\,\ln\left[1-\mathrm{e}^{-\beta\omega_\pi(\vec{q})}\right]\\ & -4\,TV\int\frac{\mathrm{d}^3p}{\left(2\pi\right)^3}\,\ln\left[1+\mathrm{e}^{-\beta(\epsilon_N(\vec{p})-\mu_B)}\right]\\ & -16\,TV\int\frac{\mathrm{d}^3p}{\left(2\pi\right)^3}\int\limits_{-\infty}^{\infty}\frac{\mathrm{d}E}{2\pi}\,\mathcal{B}\left(\vec{p},E\right)\,\ln\left[1+\mathrm{e}^{-\beta(E-\mu_B)}\right], \qquad (5)\end{aligned}$$

where the first term is the thermodynamic potential of a free pion gas and the second one is that of a free nucleon gas. The last term in eq. (5) accounts for the interaction contribution to the thermodynamic potential. It is similar to the thermodynamic potential of a free Fermi gas, but involves an integration over the energy with a weight function

$$\begin{aligned}\mathcal{B}\left(\vec{p},E\right) &= -2\,\mathrm{Im}\left[\left(1-\frac{\partial\Sigma_\Delta^R\left(\vec{p},E\right)}{\partial E}\right)\,G_\Delta^R\left(\vec{p},E\right)\right]\\ &= \mathcal{A}\left(\vec{p},E\right) + 2\,\mathrm{Im}\left[\frac{\partial\Sigma_\Delta^R\left(\vec{p},E\right)}{\partial E}\,G_\Delta^R\left(\vec{p},E\right)\right]. \qquad (6)\end{aligned}$$

---

[2] Strictly speaking we loose the nice properties of the conserving approximations in this step. However, at least for the problem at hand, the leading term in the low-density expansion is close to the self-consistent solution at moderate densities [10] ($\rho \lesssim 2\rho_0$). Self consistency is expected to be important, when there is a small scale in the problem, see e.g. ref. [11].

Here Σ_Δ^R is the retarded self energy and $\left(G_\Delta^R\right)^{-1} = \left(G_\Delta^0\right)^{-1} - \Sigma_\Delta^R$ the retarded vacuum Green function of the Δ resonance, while $\mathcal{A}$ denotes its spectral function. The functions $\mathcal{A}$ and $\mathcal{B}$ fulfill the same sum rule, i.e., the integral over all energies equals 2π.

For an interpretation of the weight function $\mathcal{B}$ in terms of diagrams it is instructive to consider the baryon density

$$n_B\left(T, \mu_B\right) = -\,\frac{1}{V}\,\frac{\partial\,\Omega\left(T, \mu_B, V\right)}{\partial \mu_B}\bigg|_{T,V}\,, \tag{7}$$

which can be split into the nucleon contribution

$$n_N\left(T, \mu_B\right) = 4\int \frac{\mathrm{d}^3 p}{(2\pi)^3}\,\frac{1}{\mathrm{e}^{\beta(\epsilon_N(\vec{p})-\mu_B)}+1} \tag{8}$$

and the contribution due to the pion-nucleon interactions

$$\tilde{n}_\Delta\left(T, \mu_B\right) = 16\int \frac{\mathrm{d}^3 p}{(2\pi)^3}\int_{-\infty}^{\infty}\frac{\mathrm{d}E}{2\pi}\,\mathcal{B}\left(\vec{p}, E\right)\,\frac{1}{\mathrm{e}^{\beta(E-\mu_B)}+1}\,. \tag{9}$$

Furthermore, as implied by the form (6) for $\mathcal{B}$, the interaction contribution (9) to n_B consists of two parts. Both of these have a straightforward interpretation: The first part, with the spectral function $\mathcal{A}$, is the contribution n_Δ of the Δ resonance to the baryon density. The second part, which is proportional to $\partial\Sigma_\Delta/\partial E$, is the contribution of the nucleons appearing in conjunction with a pion in the Δ self energy. We denote this πN pair contribution by δn_N.

We stress that there is no double counting here. The state vector of an energy eigenstate in the interacting system has amplitudes corresponding to a bare Δ and to the non-interacting πN states. The spectral function projects the interacting states onto the bare Δ state, while the second term in eq. (6) represents the projection onto the πN pair states[3]. Since the contribution of an energy eigenstate to the partition sum depends only on its energy and not on the admixture of a particular unperturbed state, both contributions in eq. (6) must be included in the thermodynamic potential, eq. (5).

One advantage of the formulation in terms of Green functions is that different contributions are easily interpreted in terms of diagrams. The three contributions to the baryon density correspond to the three diagrams shown in fig. 1(b). So far the last contribution, δn_N, has been neglected in the literature on calculations of the thermodynamics of Δ resonances [2–4]. In such calculations an *ad hoc* prescription was employed, where the spectral function $\mathcal{A}$ was used in lieu of $\mathcal{B}$.

---

[3] A detailed analysis [13] reveals that the second term not only accounts for the interacting πN pairs, but also *subtracts out* the contribution of the uncorrelated ones. This is necessary to avoid double counting, since the contribution of uncorrelated nucleons and pions is taken into account by the first two terms in eq. (5).

In order to make contact with earlier work [5,14], we show that $\mathcal{B}$ can be expressed in terms of the πN-scattering phase shift in the P_{33} channel. First, we note that the direct Δ-Born graph is the only πN-scattering process that is generated by the functional fig. 1(a). Hence, the πN-scattering amplitude is proportional to the vacuum Δ Green function G_Δ and the phase shift is given by[4]

$$\delta_{33}\left(E_{cm}\right) = \arctan\left[\frac{\mathrm{Im}\Sigma^R_\Delta\left(E_{cm}\right)}{E_{cm} - m^0_\Delta - \mathrm{Re}\Sigma^R_\Delta\left(E_{cm}\right)}\right] , \tag{10}$$

which implies that (cf. eq. (6))

$$\mathcal{B}\left(E_{cm}\right) = 2\,\frac{\partial}{\partial E_{cm}}\delta_{33}\left(E_{cm}\right) . \tag{11}$$

This form of $\mathcal{B}$ coincides with that obtained in the S-matrix formulation of statistical mechanics by Dashen, Ma and Bernstein for narrow resonances [5]. In the case we consider, namely a broad resonance, there are corrections to this expression due to interactions involving more than two particles. However, in the low-density limit two-body scattering processes dominate [15]. Thus, when the Δ self energy is properly included, eq. (4) accounts for the πN interactions in the P_{33} channel to leading order. When the logarithm in the last term of eq. (5) is expanded in powers of $\exp(-\beta(E-\mu_B))$, retaining only the lowest term, one recovers the well-known expression for the second virial coefficient of Beth and Uhlenbeck [14].

The expression (11) for the weight function $\mathcal{B}$ allows an appealing interpretation of eqs. (5) and (9). The energy derivative of the phase shift at the energy E_{cm} can be identified with the time delay of a scattered wave due to the presence of the scatterer [16]. Hence, the contribution of a resonance to the thermodynamic quantities is proportional to the time spent in the resonant state [17].

The following numerical results were obtained using the $\pi N\Delta$ interaction $\left(f_{\pi N\Delta}/m_\pi\right)\left(\vec{q}\cdot\vec{S}\right)T_a$ (cf. [18]) with a form factor of the monopole type $F(\vec{q}^{\,2}) =$

---

[4] Due to Galilean invariance the vacuum Green function and self energy depend only on the energy in the πN center-of-momentum frame E_{cm}, i.e. $\Sigma_\Delta(\vec{p},E) = \Sigma_\Delta(E_{cm})$.



Fig. 2. Upper part: Fit of δ_{33} from eq. (10) to the empirical P_{33} phase shifts [19]; Lower part: Weight function $\mathcal{B}$ (solid line), see eqs. (6, 11), compared to the spectral function $\mathcal{A}$ (dashed line).

$\Lambda^2/(\vec{q}^{\,2}+\Lambda^2)$. The parameters are chosen to reproduce the phase shift for πN scattering in the P_{33} channel [19] up to a center-of-mass energy of 1.5 GeV (upper part of fig. 2). We find $f_{\pi N\Delta} = 3.28$, $\Lambda = 290$ MeV and $m^0_\Delta = 1316$ MeV.

In the lower part of fig. 2 we compare the weight function $\mathcal{B}$ with the spectral function $\mathcal{A}$. This reveals a qualitative difference close to threshold and a strong shift of strength to lower masses, due to the second term of $\mathcal{B}$ in eq. (6). The weight function $\mathcal{B}$ is uniquely determined by the phase shifts, eq. (11), and hence a model-independent quantity. On the other hand, the spectral function and consequently the separation of $\tilde{n}_\Delta$, eq. (9), into contributions from Δ resonances and πN pairs are model dependent.

3 Pion spectra

In order to compare our calculation with experimental data we assume that at freeze-out the fireball created in a heavy-ion collision is in local thermal and chemical equilibrium. Given the population of states $\tilde{n}_\Delta$, which decay into a nucleon and a pion after freeze-out, it is straightforward to compute the resulting pion spectrum, cf. [3,20]. The pion momentum distribution in the Δ rest frame is folded with the resonance velocity distribution due to thermal motion and collective flow. The collective radial flow [4] is taken into account by assuming a flow profile, which depends quadratically on the radial coordinate[5].

The final π^0 spectrum is then a function of the temperature T, the chemical potentials for baryon number μ_B and electric charge μ_Q as well as the mean flow velocity $\bar{v}$. The parameters at freeze-out are determined by fitting the π^0 spectrum [21] of the reaction Au+Au at $\mathrm{E}_{lab}/\mathrm{A} = 1$ GeV with the constraints that the ratio of charge to baryon number equals $N_Q/N_B = 79/197$ and that the ratio of π^+ mesons to the number of participating baryons is given by [22] $N_{\pi^+_{obs}}/N_B \simeq 0.025$.

In fig. 3 we show the different contributions to the spectrum as a function of the transverse pion mass $m_\perp = \sqrt{p_\perp^2 + m_\pi^2}$ in a representation, where a Boltzmann distribution would appear as a straight line with slope $-T^{-1}$. At large $m_\perp$, the spectrum is dominated by thermal pions, while the πN pairs give an important contribution to the pion spectrum at small transverse masses. Since the weigth function $\mathcal{B}$, which determines the sum of the Δ and πN-pair contributions, is model independent, the full line is not affected by assumptions concerning the structure of the Δ resonance in vacuum, provided the phase shifts are reproduced. At freeze-out we find a temperature of $T = 57$ MeV, a baryon chemical potential of $\mu_B = 830$ MeV, a charge chemical potential of $\mu_Q = -18$ MeV and a mean flow velocity of $\bar{v} = 0.36$. The corresponding freeze-out density is $n_B = 0.4\,\rho_0$, where $\rho_0 = 0.16$ fm^{-3} denotes normal nuclear matter density.

As mentioned above the separation of $\tilde{n}_\Delta$ into a bare Δ and πN pair contribution is model dependent. Nevertheless, it is interesting to see how this is realized in a typical case. Using the spectral function $\mathcal{A}$ shown in fig. 2 the fraction of Δ resonances at freeze-out is $n_\Delta/n_B = 0.035$ while that of Δ's and πN pairs is $\tilde{n}_\Delta/n_B = 0.054$. When the Δ resonance is treated as a stable particle ($\Gamma_\Delta = 0$) one finds $n_\Delta/n_B = 0.037$ for the same values of T and μ_B.

---

[5] This choice is not crucial, because we find only a weak dependence of the final pion spectrum on the assumed analytic form of the flow profile, for a given value of the mean flow velocity.



Fig. 3. The π^0 spectrum of Au+Au at $E_{lab}/A = 1$ GeV of ref. [21]. The dashed line shows the contribution of thermal pions, while the long-dashed line corresponds to thermal pions plus those which stem from Δ resonances, decaying after freeze-out. The latter contribution is determined by the spectral function $\mathcal{A}$, fig. 2. Finally, the full line includes in addition the contribution of the πN pairs.

In this case the contribution of the correlated πN pairs vanishes.

We note that by arguments analogous to the ones presented here, one finds that the invariant mass distribution of πN pairs [23] emitted from a fireball should not be directly identified with a thermally weighted Δ resonance. Again the correlated πN pairs must be accounted for by using $\mathcal{B}$ rather than $\mathcal{A}$.

4 Summary and conclusions

We have constructed the thermodynamic potential and the baryon density of an interacting πN system taking the width of the Δ resonance consistently into account. At low densities, where medium effects can be neglected, the weight function for the Δ resonance is given by its spectral function plus a correction term. The latter accounts for the correlated πN pair component of a physical Δ resonance. The full weight function can, in agreement with the virial expansion, be expressed in a model-independent way in terms of the energy derivative of the πN scattering phase shift.

The frequently employed prescription, where the thermodynamics involves a trivial spectral function for the nucleon (δ-function) and the free spectral function of the Δ resonance, is incorrect. In such an approach, the contribution of πN pairs is neglected. These pairs yield an important contribution to the population of Δ-like states and to the low-momentum pion spectrum in relativistic heavy-ion collisions. When we include all contributions, the thermal pions, those due to Δ decays, as well as the contribution due to πN pairs, we find a good description of the experimental π^0 spectrum of Au+Au at 1 GeV per nucleon.

We thank E. Grosse, V. Metag, H. Oeschler, P. Senger, C. Sturm and A. Wagner for rewarding discussions.

References

[1] R. Stock, Phys. Rep. 135 (1986) 259
D. Hahn and N.K. Glendenning, Phys. Rev. C 37 (1988) 1053
J. Sollfrank, P. Koch and U. Heinz, Phys. Lett. B 252 (1990) 256
G.E. Brown, J. Stachel and G.M. Welke, Phys. Lett. B 253 (1991) 19
D.N. Voskresensky, Nucl. Phys. A 555 (1993) 293
P. Braun-Munzinger, J. Stachel, J.P. Wessels and N. Xu, Phys. Lett. B 344 (1995) 43

[2] D. Hahn and H. Stöcker, Nucl. Phys. A 452 (1986) 723
K.G. Denisenko and St. Mrówczyński, Phys. Rev. C 35 (1987) 1932
M.I. Gorenstein, St. Mrówczyński and D.H. Rischke, Phys. Lett. B 243 (1990) 327
J. Sollfrank, P. Koch and U. Heinz, Z. Phys. C 52 (1991) 593
R. Rapp and J. Wambach, Nucl. Phys. A 573 (1994) 626

[3] M.I. Gorenstein, M.S. Tsai and S.N. Yang, Phys. Rev. C 51 (1995) 1465

[4] P. Danielewicz, Phys. Rev. C 51 (1995) 716

[5] R. Dashen, S. Ma and H.J. Bernstein, Phys. Rev. 187 (1969) 345
R.F. Dashen and R. Rajaraman, Phys. Rev. D 10 (1974) 694
R.F. Dashen and R. Rajaraman, Phys. Rev. D 10 (1974) 708

[6] A.Z. Mekjian, Phys. Rev. C 17 (1978) 1051
M. Schmidt, G. Röpke and H. Schulz, Ann. Phys. (N.Y.) 202 (1990) 57
E.V. Shuryak, Nucl. Phys. A 533 (1991) 761
R. Venugopalan and M. Prakash, Nucl. Phys. A 546 (1992) 718

[7] G. Baym and L.P. Kadanoff, Phys. Rev. 124 (1961) 287
G. Baym, Phys. Rev. 127 (1962) 1391

[8] J.M. Luttinger and J.C. Ward, Phys. Rev. 118 (1960) 1417
A.A. Abrikosov, L.P. Gorkov and I.E. Dzyaloshinski, *Methods of quantum field theory in statistical physics*, Dover Publications, New York (1963)
R.E. Norton and J.M. Cornwall, Ann. Phys. (N.Y.) 91 (1975), 106

[9] G.M. Carneiro and C.J. Pethick, Phys. Rev. D 11 (1975), 1106
T.D. Lee and M. Margulies, Phys. Rev. D 11 (1975), 1591
E.M. Nyman and M. Rho, Nucl. Phys. A 268 (1976), 408

[10] W. Weinhold, B. Friman and W. Nörenberg (work in progress)

[11] M. Lutz, Phys. Lett. B (in press), GSI preprint 97-55, nucl-th/9709073

[12] A.F. Fetter and J.D. Walecka, *Quantum theory of many-particle systems*, McGraw-Hill, New York (1971)

[13] W. Weinhold, Diploma thesis, TH Darmstadt, 1995 (unpublished)

[14] E. Beth and G.E. Uhlenbeck, Physica 4, (1937) 915
K. Huang, *Statistical mechanics*, Wiley, New York (1963)

[15] C.B. Dover, J. Hüfner and R.M. Lemmer, Ann. Phys. (N.Y.) 66 (1971) 248

[16] E.P. Wigner, Phys. Rev. 98 (1955) 145
M.L. Goldberger and K.M. Watson, *Collision theory*, Wiley, New York (1964)

[17] P. Danielewicz and S. Pratt, Phys. Rev. C 53 (1996) 249

[18] T. Ericson and W. Weise, *Pions and Nuclei*, Clarendon Press, Oxford (1988)

[19] R.A. Arndt, I.I. Strakovsky, R.L. Workman and M.M. Pavan, Phys. Rev. C 52 (1995) 2246

[20] R. Hagedorn, *Relativistic Kinematics*, Benjamin, Reading (1963)

[21] O. Schwalb et al., Phys. Lett. B 321 (1994) 20

[22] P. Senger, Acta. Phys. Pol. B 27 (1996) 2993

[23] E.L. Hjort *et al.*, Phys. Rev. Lett. 79 (1997) 4345